NON-TUBERCULOUS MYCOBACTERIUM: MANAGEMENT

Doreen J. Addrizzo-Harris,MD

Professor of Medicine

Co-Director, NYU Bronchiectasis Program

Associate Director for Education and

Faculty Affairs

Division of Pulmonary, Critical Care and Sleep Medicine

- Nutrition
  -

NYU Langone Health

MAC fibronodular



MAC Fibronodular Lung Disease: Treatment

- 180 patients
  - Completed > 12 Months of macrolide/azalide therapy
  - No differences in clarithromycin vs azithromycin
  - Treatment modification increased in daily (80%) vs intermittent (1%)
  - Treatment success achieved in 84% of patients
  - Microbiologic recurrences in 48% after completion of therapy – 75% reinfection isolates/25% true relapse
    - » Wallace RJ, et al Chest 2014 Aug 146:276-82

- Primary endpoint was change from baseline to day 84 on a semi-quantitative mycobacterial growth scale.
    - Other endpoints included sputum conversion, 6-minute walk distance, and adverse events.
- Primary endpoint was not achieved (P=0.072); however, a greater proportion of the LAI group demonstrated ≥1 negative sputum cultures (32% [14/44] vs. 9% [4/45]; P=0.006) and improvement in 6-minute walk test (+20.6 vs. −25.0 meters; P=0.017) at day 84.

NYU Langone Health

Randomized Trial of Liposomal Amikacin for Inhalation in Nontuberculous Mycobacterial Lung Disease

Olivier, et al Am J Resp Crit Care Med 2017

Serious events

- In the double-blind phase, the overall incidence of serious adverse events was higher in the LAI group compared with the placebo group (18.2% vs. 8.9%)
  - bronchiectasis exacerbation (2 LAI; 1 placebo)
  - pneumonia (1 LAI; 2 placebo)

- Renal events – 1 mild event; Audiology - 5 events

NYU Langone Health

Conclusions

- This study indicates that in select patients with treatment-refractory non-CF MAI disease, LAI added to guidelines-based therapy can achieve early and sustained negative sputum cultures.
-

NYU Langone Health

- Results
  - Ethambutol

NYU Langone Health

Treatment duration and monitoring

- Treatment is usually 18-24 months;
- A minimum of 12 months after culture conversion

- Monitor patient closely for response and for side effects; need baseline and serial:
  - Labs
  - EKG
  - Audiometry
  - Ophthalmology exam
  - Physical exam and weights
  - Monthly sputums for AFB
  - Follow up CT scan/ PFTs

NYU Langone Health